



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 15/29, 15/63, 15/67 C12N 15/82, A01H 4/00, 5/00	A1	(11) International Publication Number: WO 91/09948 (43) International Publication Date: 11 July 1991 (11.07.91)
(21) International Application Number: PCT/US91/00073 (22) International Filing Date: 4 January 1991 (04.01.91) (30) Priority data: 461,490 5 January 1990 (05.01.90) US (71) Applicant: CORNELL RESEARCH FOUNDATION, INC. [US/US]; 20 Thornwood Drive, Ithaca, NY 14850 (US). (72) Inventors: WU, Ray ; 111 Christopher Circle, Ithaca, NY 14850 (US). McELROY, David ; 96 Lowercreek Road, Etna, NY 13062 (US). (74) Agent: YAHWAK, George, M.; Yahwak & Associates, 25 Skytop Drive, Trumbull, CN 06611 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), SE (European patent), SU. Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: RICE ACTIN GENE AND PROMOTER (57) Abstract The present invention describes the genomic nucleotide sequence, and isolation of a strong intron-dependent promoter from rice actin genes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland			SN	Senegal
CI	Côte d'Ivoire	KR	Republic of Korea	SU	Soviet Union
CM	Cameroon	LI	Liechtenstein	TD	Chad
CS	Czechoslovakia	LK	Sri Lanka	TG	Togo
DE	Germany	LU	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		

RICE ACTIN GENE AND PROMOTER

Significant progress has recently been made in the application of tissue culture and gene transfer techniques to previously recalcitrant monocotyledonous plants, such as rice. Rice protoplasts have been shown to transiently express a foreign gene after transformation by electroporation, and stable genomic integration of foreign DNA, following polyethylene-glycol-mediated transformation of rice cells, has been previously demonstrated.

One of the major limitations in rice transformation technology has been the lack of an efficient promoter for high level constitutive expression of foreign genes in transgenic plants. The cauliflower mosaic virus (CaMV) 35S promoter has been widely used for this purpose in a number of plant transformation systems, however, the CaMV 35S promoter has shown low activity in transforming rice cells, and recent reports suggest that the pattern of CaMV 35S promoter activity in transgenic plants may not be constitutive. Work in our laboratory suggests that the 5' region of the maize Adh1 gene containing the promoter, 5'-coding exon and 5'-intron, is 20 to 30 times more active than the CaMV 35S promoter for transient gene expression in transformed rice protoplasts and calli, however, the maximum activity of AdH1 promoter in transformed rice protoplasts requires anaerobic stress and its activity is not constitutive in all transformed rice tissues. As a result, a number of alternative

promoter elements for rice transformation are currently under investigation.

It is, therefore, the purpose of the present invention to describe an efficient promoter for rice transformation, and in
5 doing so have based such a promoter upon a rice actin gene that displays a constitutive spatial and temporal pattern of transcript abundance.

Cytoplasmic actin is a fundamental component of the eukaryotic cell cytoskeleton. In higher-plant cells a number of
10 cellular process, such as cytoplasmic streaming, extension growth and cell division, are believed to involve cytoskeletal actin protein. Actin has been found to be encoded by gene families in all higher plants studied to date. A unique feature of these plant actin genes, in contrast to the situation for animal
15 actin genes, has been their conservation of gene structure. All of the studied plant actin genes are reported to consist of four coding exons of conserved length, separated by three introns of variable length. In each of these cases the coding regions of the plant actin genes have been deduced following a comparison of
20 their potential translations products with that of previously published animal actin protein sequences. However, in a number of animal actin genes there exists a 5' transcribed, but untranslated, sequence (5'-noncoding exon), separated by an intron (5'-intron) from the exon containing the first translated codon
25 (5'-coding exon). Although it has been suggested that a 5'-noncoding exon may exist in plant actin genes, such an exon would

fail to be detected in any comparison of the coding region of plant and animal actin genes.

In rice, there are at least 8 actin-like sequences per haploid genome, of which four have been isolated and shown to differ from each other in the tissue and stage specific abundance of their respective transcripts. One rice actin gene, RAc1, was found to encode a transcript that is relatively abundant in all rice tissues, and at all developmental stages examined. In view of the high level constitutive abundance of the rice RAc1 transcript the present invention describes the isolation and partial sequencing of a genomic clone containing the rice RAc1 5' flanking and 5'-transcribed, non-translated, regions. From such clones, a number of RAc1-GUS (GUS stands for β -glucuronidase gene) fusions plasmids were constructed and used in transient expression assays of transformed rice protoplasts. The results from these assays suggest that the regulatory element(s) necessary for maximal RAc1 promoter activity in transformed rice protoplasts are located within a region 1.3 kb upstream of the RAc1 translation initiation codon. The rice RAc1 promoter is 5 times more active than the maize Adh1 promoter in transformed rice cells, suggesting that the RAc1 5' region contains an efficient promoter for rice transformation.

As a first step towards a characterization of the actin gene in rice, several clones from a rice genomic library representing four unique actin sequences were isolated. A rice (*Oryza sativa* variety IR26) genomic library in lambda-EMBL4 was screened

with a heterologous actin probe. Fourteen independent clones were isolated and subcloned into pUC13. By mapping restriction sites and carrying out cross-hybridization studies, four different classes of clones were identified and designated RAc1, RAc2, 5 RAc3 and RAc7.

Sequencing of the subcloned actin genes was carried out by the dideoxynucleotide chain termination method and computer analysis of the resulting sequences was done. The nucleotide sequences of the coding regions of the four actin genes appear in 10 the EMBL Gene Bank and DDB7 Nucleotide Sequence Databases under the accession numbers X15865 RAc1; X15864 RAc2; X15862 RAc3; and X15863 RAc7.

Prior to identifying those sequences which regulate RAc1 expression, its complete genomic structure was determined. By 15 characterizing an RAc1 cDNA clone (EMBL data bank accession number X16280) and 5' end mapping the RAc1 transcript, the structure of the RAc1 gene was determined and the position of a 5'-noncoding exon in its genomic sequence was identified and located. This represents the first complete structural 20 characterization of plant actin gene and is one of few reported cases of a 5'-noncoding exon in a plant gene.

A complete understanding of all aspects of the present invention can be better understood from reference to the accompanying figures (and examples) in which:

25 Figure 1a is a restriction map of the pcRAc1.3 cDNA insert, according to the present invention;

Figure 1b is a restriction map and proposed structure of the RAc1 genomic clone according to the present invention;

Figure 2a is a restriction map of the 15.1 Kb lambda-RAc1 insert, according to the present invention;

5 Figure 2b is a restriction map of the pRAc 15'.H3 insert according to the present invention;

Figure 3 depicts maps of constructs according to the present invention.

The identification of the complete genomic structure was
10 accomplished utilizing the protocols described in the following examples:

Example 1

a. Screening the lambda-gt11 c DNA library:

- A rice (*Oryza sativa* IR36) lambda-gt11 cDNA library,
5 prepared from six-day-old etiolated leaf tissue was plated and
lifted onto nylon filters. Primary and secondary screenings for
rice actin cDNA clones were carried out by hybridization with a
260 bp, ³²P-dATP labeled BamH1-HindIII restriction fragment
from the rice actin genomic clone pRAc2. Tertiary screening was
10 carried out using rice actin gene- specific probes. RAc1-like
cDNA clones were identified using a 900-bp BamH1-HindIII
restriction fragment probe from the 3' end of the rice actin
genomic clone pRAc1.

15 b. Restriction mapping and nucleotide sequencing:

- Actin positive clones from tertiary screening of the
lambda-gt11 cDNA library were further characterized by
restriction mapping. Restriction fragments from the cDNA insert
of the RAc1 positive clone pcRAc1.3 were subcloned into pBS-KS
20 vectors for nucleotide sequencing. Double-stranded DNA
sequencing using ³⁵S-dATP was carried out using a T7
polymerase sequencing kit following the manufacturer's
(Pharmacia) modification of the Sanger *et al* dideoxynucleotide
sequencing procedure. The 5' and 3' ends of the pcRAc1.3 cDNA
25 insert were further sequenced using two synthetic
oligonucleotides; 5' -AAGCGGCCTCTACTACGC-3' and 5'-

GAAGCATTTCTGRCACAAT-3' respectively and subsequently the sequence data were analyzed.

- 5 c. 5'-end mapping of RAc1 transcript by northern blot analysis and primer extension:

10 RNA isolation from seven-day-old rice shoots and Northern blotting were performed. 10 µg of total RNA samples were used in formaldehyde agarose gel electrophoresis and northern analysis was carried out under stringent hybridization conditioned (50% formamide, 50°C). Restriction fragment probes for northern hybridization were isolated from the 5'-untranslated end of the rice actin genomic clone pRAc1 and subcloned into pBS-KS vectors.

- 15 The primer extension analysis was performed using 1 µg of a synthetic oligonucleotide primer 5'-CTTCTACCGCGGCGGC-3' which was annealed to 25 µg of total RNA from seven-day-old rice shoots.

The cDNA library made with mRNA from six-day-old etiolated rice shoots according to Example 1 provided the cDNA clones corresponding to the multiple members of the rice actin gene family. Primary and secondary screening with a 260 bp *Bam* H1-*Hind* III restriction fragment - a fragment previously known to cross-hybridize with many members of the rice actin gene family - from the rice actin gene RAc2, because of the homology found among isolated rice actin genomic sequenced. Thirty four clones were identified which strongly hybridized to the non-specific RAc2 actin probe.

Tertiary screening of the 34 actin-positive cloned was carried out using probes previously determined to be actin gene specific. Seven RAc1-like cDNA clones were identified which hybridized specifically to a 900 bp *Bam*H1-*Hind*III restriction fragment from pRAc1. The nature of these clones was confirmed by restriction mapping and Southern blotting. The *Eco*RI insert from the lambda-gt 11 clone lambda- RAc1.3 was subcloned into a pBS-KS vector to produce the plasmid pcRAc1.3.

Using similar procedures, nine RAc3-like cDNA clones and eight RAc7-like cDNA clones were identified. No cDNA clones were identified which cross-hybridized with a RAc2 gene specific probe. The remaining ten actin-positive clones failed to cross-hybridize with any of the previously characterized actin gene-specific probes. These were classified into five groups on the basis of restriction mapping and cross-hybridization analysis. The isolation of cDNA clones representing eight distinct actin

transcripts confirms that the actin gene family in rice composed of at least eight unique members.

The insert from pcRA1.3 was further subcloned into pBS-KS plasmids. Double stranded DNA was prepared for sequencing. The
5 restriction enzyme map of the pcRAC1.3 insert and the sequencing strategy for determining its base are summarized in Fig. 1a.

More specifically, Fig. 1 describes the determination of the RAC1 gene structure by alignment of the pcRAC1.3 cDNA insert with the pRAC1 genomic clone a: restriction map of the pcRAC1
10 cDNA insert with the strategy used to determine its base sequence. The horizontal arrows indicate the direction of sequencing and their length indicate the actual size of the sequence obtained. Horizontal arrows preceded with a sinusoidal wave indicate the use of synthetic oligonucleotide primers. b:
15 restriction map and proposed structure of the RAC1 genomic clone. Horizontal lines represent introns and boxes represent exons. Open boxes represent the transcribed but untranslated regions of exons, closed boxes represent the translated regions of exons. Restrictions sites: B, BamH1; H, HindIII; R, EcoRV; S, SstI.
20 Dashed lined between the cDNA and genomic DNA restriction maps indicate the alignment of their homologous exons.

The tabulated sequence which depicts the nucleotide sequence of the rice actin gene RAC1, is depicted in the following sequence of 5510 nucleotides

10

	1	GATATCCCTC	AGCCGCCCTT	CACTATCTTT	TTTGCCCGAG
		TCATTGTCAT	GTGAACCTTG	GCATGTATAA	TCGGTGAATT
	81	GCGTCGATTT	TCCTCTTATA	GGTGGGCCAA	TGAATCOGTG
		TGATCGCGTC	TGATTGGCTA	GAGATATGTT	TCCTCCTTGT
5	161	TGGATGTATT	TTCATACATA	ATCATATGCA	TACAAATATT
		TCATTACACT	TTATTAGAGG	TGGTCAGTAA	TAAACCTAT
	241	CACTATGTCT	GGTGTTTCAT	TTTATTTGCT	TTTAAACGAA
		ATTGACTTCT	GATTCAATAT	TTAAGGATCG	CAATCGCGGT
	321	GCAGTACTAA	TTCTGGTTGA	CGGAACATA	CGTAAACTAT
10		TCAGCTTCAC	TCTATTAGGC	ACTACTTGAC	ACACCGGAGT
	401	ACGTACGAAT	ACTCTATCAA	TTAGTCTCAG	TCTAACTTGT
		TGAGACATGT	ACTATAGATT	ACTATTGTAC	CTTGACTCAC
	481	TGTATGTATC	ACGTCTAATT	GAACACACA	TATATACGCG
		ATATTTTTTA	ATAACATTAA	AACCTACCTC	TATGTCAACA
15	561	ATGGTGTACG	ATAACCCAAA	GTTTAGGAGG	TAAAAAACA
		TTGCCCTACG	CTGAAGTTAC	GCCTTAAAAA	TAAAGAGTAA
	641	ATTTTACTTT	GACCACCCCT	CAATGTTTAC	TTTAGACCGG
		TGGAACGCTC	CAGCCGTAAT	AGGATTCCTG	ACCTCACATG
	721	CCTTAGCCGG	ATTATATTGC	CTGCCACCTT	TCTCACTCAT
20		ATCTGCAAGA	ATGTCTACTC	GCTAGAATTA	TCGGATAGT
	801	AGCTAGCATA	CTGAGGTCAC	TTCATATGCT	TGAGAAGAGA
		GTCGGGATAG	TCCAAAATAA	AACAAAGGTA	AGATTACCTG
	881	GTCAAAAGTG	AAAACATCAG	TTAAAAGGTG	GTATAAGTAA
		AATATCGGTA	ATAAAAGGTG	GGCCAAAGTG	AAATTTACTC
25	961	TTTTCTACTA	TTATAAAAT	TGAGGATGTT	TTGTCGGTAC
		TTTGATACGT	CATTTTTGTA	TGAATTGGTT	TTAAGTTTAA
	1041	TTGCGGATTT	TGGAATGCA	TATCTGTATT	TGAGTCGGGT
		TTTAAGTTGG	TTTGCTTTTG	TAAATACAGA	GGGATTTGTA
	1121	TAAGAAATAT	CTTTAAAAAA	ACCATATGCG	TAATTTGACA
30		TAATTTTTGA	GAAAAATATA	TATTCAGGGG	AATTCTCACA
	1201	ATGAACAATA	ATAAGATTAA	AATAGCTTGC	CCCGGTTGCA
		GCGATGGGTA	TTTTTCTAG	TAAAAATAAA	GATAAACCTA
	1281	GACTCAAAAC	ATTTACAAAA	ACACCCCTCA	AAGTCCTAAA
		GGCCAAAGTG	CTATGCAACA	TCCATAGCAA	GGCCAGCCCA

	1361	ADCCAAACCA	ADCCAAACCA	CCCCAGTGCA	GCCAACTGGC
		AAATAGTCTC	CACACCCCGG	CACATATCAC	GTGAGTTGTC
	1441	CGCACCAACG	CAGCTCTCGC	AGCCAAAAAA	AAAAAAGAA
		AGAAAAAATA	GAAAAAGAAA	AAACAGCAGG	TGGGTCCGGG
5	1521	TGTGTGGGGC	CGGAAAGCG	AGGAGGATGG	CGAGCAGCGA
		CGAGGCGGGC	CCTCCCTCCG	CTTCCAAAGA	AACGCCGCCG
	1601	ATCGCCACTA	TATACATAAC	CCCCCTCTTC	CTCCGATCCC
		CCCAACCCTA	CCACACCCAC	CACACCCACC	TCTCCGCCCG
	1681	TGCGTGGCGG	ACGAGGAGCT	CCTCCGCCCT	CCCCCTGGGC
10		CGCGCGCGGT	AACCAACCGG	CCCCCTCTCT	CTTTCTTTCT
	1761	CGGTTTTTTT	TTTCTCTCCG	GTCTCGATCT	TTGGCCTTGG
		TAGTTTGGGT	GGCGGAGAGC	GGCTTCGTGG	CCGAGATCGG
	1841	TGGCGGGGAG	GGCGGGGATC	TGGCGGCTGG	CGTCTCCGGG
		CGTGAGTCCG	CCCGGATCCT	CGCGGGGAAT	GGGGCTCTCG
15	1921	GATGTAGATC	TTCTTTCTTT	CTTCTTTTGG	TGGGTAGAAT
		TTGAATCCCT	CAGCATTGTT	CATCGGTAGT	TTTTCTTTTC
	2001	ATGATTTGTG	ACAAATGCAG	CCTCGTCCGG	AGCTTTTTTG
		TAGGTAGAAG	ATGGCTGACG	CCGAGGATAT	CCAGCCCCCT
	2081	GTCTGCGATA	ATGGAACTGG	TATGGTCAAG	GTAAGCTGTT
20		TGGATCTCAG	GGTGGTTTCC	GTTTACCGAA	ATGCTGCATT
	2161	TCTTGGTAGC	AAAACGAGGG	TGGTTTGTGT	CAGGCTGGGT
		TGGCGGGAGA	TGATGGGCCG	AGGGCTGTCT	TCCCCAGCAT
	2241	TGTGGCGCGC	CCTCGCCACA	CCGGTGTTCAT	GGTGGGAATG
		GGCCGAGAGG	ACGCCCTAGT	CGCGGAGGAG	GGCGAGTCCA
25	2321	AGAGGGGTAT	CTTGACCCCTC	AAGTACCCCA	TGAGCATGGG
		TATCGTCAGC	AACTGGGATG	ATATGGAGAA	GATCTGGCAT
	2401	CACACCTTCT	ACAAAGAGCT	CCGTGTGGCC	CCGGAGGAGC
		ACCCCGTCTT	CCTCACCGAG	GCTCCTCTCA	ACCCCAAGGC
	2481	CAATCGTGAG	AAGATGACCC	AGATCATGTT	TGAGACCTTC
		AACACCCCTG	CTATGTACGT	CGCATCCAG	GGCGTCTCTT
30	2561	CTCTGTATGC	CAGTGGTGGT	ADCCAAAGGTG	AGCACAATGG
		ACACTGAACCT	AAAAGGCTGT	GAGGATGAAT	TTTAATTTTG
	2641	ACATTTCACAT	GTAGATGAGA	TTTAGTCTTG	CAATCTTCAA
		TTGTATACAA	GCAAGACTAT	ATAATGACTT	TCAAAATAAA

5	2721	ATCATAGGCA	GTTCTCATAA	ATGGAATCAT	GTTTGAACAT
		CCTAATTCTG	TTGGCATGGA	GTGCTTTGAC	ATTTTGAAGT
	2801	GTGCATCAGT	GTGAATAACT	GAATTTCCCT	TTCCCAAGGT
		ATTGTGTGGG	ACTCTGGTGA	TGGTGTGAGC	CACACTGTCC
10	2881	CCATCTATGA	AGGATATGCT	CTCCCCCATG	CTATCTCTCG
		TCTCGACCTT	GCTGGGCGTG	ATCTCACTGA	TTACCTCATG
	2961	AAGATCCTGA	CGGAGGCTGG	TTACTCATTG	ACCACAACTG
		CCGAGCGGGA	AATTGTGAGG	GACATGAAGG	AGAAGCTTTT
15	3041	CTACATGGCC	CTGGAATATG	ACCAGGAAAT	GGAGACTGCC
		AAGACCACTG	CCTCGGTGGA	GAGGAGCTAC	GAGCTTCTCG
	3121	ATGGACAGGT	TATCAACATT	GGTGTCTGAG	GTTTCCGCTG
		CCCTGAGGTC	CTCTTCCAGC	CTTCTTCCAT	AGGAATGGAA
20	3201	GCTGGGGTGA	TCCATGAGAC	TACATACAAC	TCCATCATGA
		AGTGGGAGCT	GGATATTAGG	AAGGATCTAT	ATGGCAACAT
	3281	CGTTCTCAGT	GGTGGTACCA	CTATGTTCCG	TGGCATTTGT
		GACAGGATGA	GCAAGGAAGA	TCACTGCCCT	GCTCTAGCCA
25	3361	GCATGAAGAT	CAAGGTGGTC	GCCCCCTCTG	AAAGGAAGTA
		CAGTGTCTGG	ATTGGAGGAT	CCATCTTGGC	ATCTCTCAGC
	3441	ACATTCCAGC	AGGTAATAT	ACAAATGCAG	CAATGTAGTG
		TTGTTTACCT	CATGAACCTG	ATCAATTTGC	TTACAATGTT
30	3521	GCTTGCCGTT	GCAGATGTGG	ATTGCCAAGG	CTGAGTAAGA
		CGAGTCTGGC	CCATCCATTG	TGCACAGGAA	ATGCTTCTAA
	3601	TTCTTCGGAC	CCAAGAATGC	TAAGCCAAGA	GGAGCTGTTA
		TGGCCGTCTT	CCTGCTTGTT	TCTCTCTTTT	TGTTGCTGTT
35	3681	TCTTCATTAG	CGTGGACAAA	GTTTTCACCC	GCCCTATCTG
		TTATCATTTT	CTTCTATTCA	AAGACTGTAA	TACCTATTGC
	3761	TACCTGTGGT	TCTCACTTGT	GATTTTGGAC	ACATATGTTT
		GGTTTATTCA	AATTTAATCA	GATGCTGAT	GAGGGTACCA
40	3841	GAATAAATAC	GTGTTCTGGT	TGTTTTTGAG	TTGGGATTAT
		TCTATGAAAT	GAATAACATC	GAAGTTATCA	TCCAGTATT
	3921	TTGGCATGAA	TGTTCTTTTC	TTCTGTCTTG	TGCATCAGTG
		ATCTAGTGCA	TGGGAGTTTG	TATTGTGATG	TTGCACATCA
45	4001	CGTAACCTCC	ACTTTGCCCT	TGCTGTTCGA	TATTTTAATG
		ACATGTCCAA	CACACTTCTG	CTACTTTTCT	TTCTTGGCTA

	4081	TTGTGCGAGC	ATGATGCAAG	ATGCATCACA	CGATCAGATA
		TATTCTCATC	GTCAGGCTTT	CAGGCGACGA	GCACGCTTTG
	4161	CGCTTAAAG	TTGTACCGGC	AGTAGACATC	CCCTGTAGAA
		GTGATAATCT	TTTCACTTTT	CTTAAAGAAA	TTGAGAGGGG
5	4241	AAATGGAACC	ATGTGGATCA	GAGAAGCTTT	TGTTTCTTAC
		ACAAGAATAT	TTGGTACAGT	GGGGGTCTTA	TGTTGCTGGG
	4321	TTGGTGGCTT	GGCTCCCTGT	CTTCAACCAA	GTGTTTTGAG
		TTCAACATGT	TAGCGTGTAG	AAAGAGCACA	ATTCTGTTTA
	4401	TCTCAAGGT	AAAATGTGGC	ATTCTGTTAA	AGAACATGAT
10		CCTGCGAATT	TTTTAAGTTT	CAATGGAAGA	GGAATGTAAA
	4481	GCTTTCTATG	GTTTGTGTAC	ACAACACAGT	GGAAGAGGAG
		TGCAAGCTTT	CTATGGTTTG	TGTGCGGCTT	GTGGTGCAGC
	4561	ACTTCAATTT	TGTTAGAAAT	GAAGAAGAAA	AAAGATGAT
		CATGCTTATA	GTAATCACT	CTTTTTCTCT	GCCTCTGTGA
15	4641	CGTTTTGACT	TGACAAGATT	TTAAATCTGT	TACATGAOCT
		TTGTTTTAAA	ATTACTTTAT	GTATTTCCAT	CTTTCAAGTT
	4721	ATGCAGATGT	CATCACAAAT	TGTTACACCA	ATCACAGGCC
		TGGCTGTTTTA	TATATTATCA	GAOCAGGCTA	TATAGAGTAT
	4801	ACTATACTAA	CTGTTCATAT	TATCTGGAAG	TCTTGCCTGC
20		TACTTGAGCG	GGAAAGGGT	ATAGATATGA	GGTTAAGGAA
	4881	CGAAGCGCA	GCAATCGAG	GCTCTCTCTG	AAATCATTTT
		ACATCTACAA	AAGCACATTT	AACCTTTTCT	AGAACACATA
	4961	TGTTACTTAG	AAGCAGGAAG	TTTATGCAAA	ATTTTCATCGA
		CAAGATAACC	AGGGGGGCAC	TGGAAGAGTT	ATCTTTTACC
25	5041	TCAATCTGTA	TACACTCAAA	GTTACTCGGA	TTGTACATTG
		GCTAAAAGTT	TCCCTGTTTC	ATTTGAACCA	CCTCAGCAAA
	5121	AGCAACCTGA	AGAGTTTGTT	GTGCAAGGT	AAAAACCTTC
		CCCCAGACTT	TGATCCTTCC	CTTGCAATATC	TAAGGGCATC
	5201	AOGGTGAGGT	CACGTATACG	CAAGCATTAG	TCCAACACAA
		AGCCATTCTT	TGCTTCTTTT	GTCCACCGTT	TCAATATGTA
30	5281	TACATCTGGT	ATGTTGCGTA	CATCAAGGGC	CAAGAATACT
		CTTAGTATAT	GGCGGCACAA	GCTACCACAA	CTCTCAAACT
	5361	TGCAGCAGCT	GCACTTAGCT	ATATTGCCAG	AAGTATCATA
		CCTGACTCTG	CATGTGGCTT	CAGTATGGTC	CTTTGTGACA

5441 CTATACACAG CAATCAACCC ATCATTGTCA AGACTAGAGA
TATATAATAG CCTAAAGATC CAATGAATCC 5510

An alignment between the sequence of the RAc1 cDNA clone and that of the RAc1 genomic sequence was used to determine the structure of the RAc1 gene shown above. Translation of the pcRA1.3 insert in all three reading frames identified a potential coding region of 1131 nucleotides. The potential coding region of the RAc1 cDNA, if translated *in vivo*, would code for 377 amino acids and an actin protein of 41.9 kDA estimated molecular mass. This analysis identified three introns, interrupting the gene at the same places in the RAc1 coding sequence as those previously reported for all other plant actins. The analysis also identified an additional intron within the transcribed sequence of RAc1 which is 5' of the region containing the translation initiation codon. This 5'-intron separated a 79 bp GC-rich 5'-noncoding exon from an exon coding the translation initiation codon.

In the sequence depicted above, the promoter according to the present invention lies within 0 to 2071 nucleotides, more particularly, the efficiency of the promoter appears to lie within about 800 to 2071 nucleotides; nucleotides 811 to 816 and 2066 to 2070 are the XhoI and EcoRV enzyme sites, respectively; 2051 to 3600 nucleotides encompass the coding region DNA sequence for the rice actin gene RAc1; and 1650 to 3841 nucleotides is the RAc1 genomic clone for pRAc1.

To determine if any of the previously isolated plant actin genes also contain 5'-intron-like sequences, the region 5' of their

respective translation initiation codons to that of RAc1, and those animal actin genes known to contain such 5'-introns as compared. This analysis revealed that the soybean (*Glycine max*) actin genes SAc3 and SAc1, the *Arabidopsis thaliana* actin gene AAc1, the potato (*Solanum tuberosum*) actin processed pseudogene PAC-psi, and the maize (*Zea mays*) actin gene MAC1 have regions upstream of their translation initiation codons which bear sequence similarity to the 3'-splice site junction regions of the 5'-introns of the rice actin gene RAc1, a *Xenopus borealis* cytoplasmic actin gene and the *Drosophila melanogaster* cytoplasmic actin gene DmA2. These previously confirmed, and putative intron splice sites are all found within 7 to 11 bp upstream of their respective translation initiation codons. Although the short region of untranslated exons they would encode are all AT-rich, they do not as a group suggest any strong consensus sequence. However, these sequences may serve a similar function in the different actin genes.

The complete structural analysis of the RAc1 gene described above has therefore led to the identification and localization of a 5' noncoding exon, separated by a 5'-intron from the first coding exon, in the RAc1 genomic sequence. It has been reported that a 5'-intron in the maize Adh1 gene is essential for the efficient expression of foreign genes from the maize Adh1 promoter [Callis et al., 'Introns increase gene expression in cultured maize cells. *Genes & Development*, 1:1183 (1987)]. In order to investigate the effect of the RAc1 5'-intron on gene

expression, a number of GUS fusion plasmids containing RAc1 intron deletions which involved the successive removal of those intron sequence elements previously determined as being important for efficient intron splicing, such as the mRNA branch point and 3'-donor splice sites. The results of transient assays of GUS activity in rice protoplasts transformed with the various deletion constructs suggest that the 5'-intron of RAc1 is essential for efficient gene expression from the RAc1 promoter. Test results also suggest that the intron-mediated stimulation of gene expression is not a function of the intron sequence per se but is associated, in part, with an in vivo requirement for efficient intron splicing.

This analysis and following discussion of the identification and localization of the 5' noncoding exon in the RAc1 genomic sequence can be more easily understood with references to the following examples:

Example II

Genomic Clone Characterization:

- A genomic restriction map of the 15.1 kb insert from
- 5 lambda-RAc1 was prepared by analyzing all possible single and double digests with the enzymes BamHI, EcoRI, HindIII and Sall. A 5.3 kb HindIII-HindIII restriction fragment from the lambda-RAc1 clone was subcloned into pBluescript-KS to generate the plasmid pRAc15'.H3. A restriction map of pRAc15'.H3 was prepared by
- 10 analyzing all possible single and double digests with the enzymes BamHI, BglII, EcoRI, EcoRV, HincII, HindIII, KpnI, PstI, SmaI, SphI, XbaI and XhoI. Restriction digestion, plasmid ligation, transformation of E. coli DH5-alpha competent cells and isolation of plasmid DNA were done following standard procedures.
- 15 A 2.1 kb EcoRV-EcoRV region within the pRAc15'.H3 insert was further subcloned into pBluescript-KS and its DNA sequence determined according to the sequencing strategy outlined in Fig. 2B. Double stranded DNA sequencing reactions were run using the dideoxy chain termination method following T7 polymerase™
- 20 protocol. Analysis of DNA sequence data were performed using the Microgenie™ computer program.

EXAMPLE III

Construction of Plasmids:

The 2.1 kb EcoRV-EcoRV fragment from pRac15'.H3 was
5 subcloned into the SmaI site of pBluescript-KS, in both
orientations, to produce the plasmids pRac15'.21V and
pRac15'.21V^r. A promoterless β -glucuronidase (GUS) gene,
containing the 3' noncoding region of the nopaline synthase (NOS)
gene, was excised by BamHI-XbaI digestion of pEXAG3 and cloned
10 between the BamHI and XbaI sites of pRac15'.21V and
pRac15'.21V^r to produce the plasmids pRac15'.21VG and
pRac15'.21V^rG. Deletion of the 0.8 kb XhoI-XhoI and 1.2 kb EcoRI-
EcoRI fragments from pRac15'.21VG generated the plasmids
pRac15'.13XG and pRac15'.09RG respectively. Introduction of the
15 1.2 kb EcoRI-EcoRI fragment into the EcoRI site of pRac15'.09RG,
in the reverse orientation, produced the plasmid pRac15'.21V^rG.
The plasmid pRac15'.09RGIDS⁻ was produced by deletion of a 0.17
kb BamHI-BamHI fragment from pRac15'.09RG.

The plasmids pRac15'.09RGI Δ 8, pRac15'.09RGI Δ 1 and
20 pRac15'.09RGI Δ 12 were generated by cleavage of pRac15'.09RG at
its BglII site followed by exonuclease III deletion and S1 nuclease
treatment to remove different amounts of the RAC1 first intron.
To construct the plasmid pRac15'.09RGI⁻ the 0.9 kb EcoRI-EcoRV
fragment from pRac15'.H3, containing the first intron of the rice
25 RAC1 gene, was cloned between the EcoRI and EcoRV sites of
pBR322 to produce the plasmid pBRac15'.09R. The 0.4 kb SstI-

EcoRV fragment from pBRAc15'.09R, containing the RAc1 intron, was excised and replaced with the intronless 0.1 kb SstI-EcoRV fragment from the insert of an RAc1 cDNA clone, pcRAc1.3, to produce the plasmid pBRAc15'.09RI⁻. The intronless 0.6 kb EcoRV
5 fragment from pBRAc15'.09RI⁻ was excised and cloned between the EcoRI and SmaI sites of pBS-KS to yield the plasmid pRAc15'.09RI⁻. Cloning of the Gus-Nos containing BamHI-XbaI fragment from pEXAG3 between the BamHI and XbaI site of pRAc15'.09RI⁻ produced the plasmid pRAc15'.09RGI⁻. The nature
10 of this RAc1 promoter-GUS fusion was confirmed by sequencing double-stranded DNA.

EXAMPLE IV

Culture, transformation and visualization of GUS activity in rice cell suspension cultures and protoplasts:

- 5 Cell suspension cultures were generated from calli, initiated from scutella of mature rice (*Oryza sativa* L. v Lemont) seeds, and cultured in liquid R2 media containing 3% (w/v) sucrose, 2 mg/ml 2,4-dichlorophenoxyacetic acid and 2 mg/l vitamin B-5. All cell suspensions were subcultured weekly and
10 incubated in the dark at 26°C. Cell suspensions were filtered through a 700 µm mesh prior to particle gun bombardment with 1.2 µm diameter tungsten particles mixed with a solution containing 10 µg of plasmid DNA. GUS activity in intact cells and developing calli was determined by the appearance of blue spots
15 two and ten days after bombardment with plasmid DNA following the GUS assay procedure of Jefferson et al.

- Protoplasts were isolated from rice (*Oryza sativa* L. v Nipponbare) cell suspension cultures and resuspended in MaMg medium to a final density of 10^6 protoplasts/ml. For
20 transformation, 1 ml of protoplast suspension was incubated with 10 µg of circular plasmid DNA, 50 µg/ml of calf thymus carrier DNA and an equal volume of 30% polyethylene glycol 4000. The mixture was incubated for 30 minutes, diluted with CPW13 medium, washed a further 3 times in CPW13 media, with
25 centrifugation between each wash, before being resuspended in simplified KPR liquid media to a final density of 10^6

protoplasts/ml. 0.1 ml aliquots of this final suspension were incubated by thin layer culture in 96-well plates prior to subsequent assays for GUS activity.

EXAMPLE V

Analysis of GUS activity in transformed protoplasts:

- To monitor the temporal pattern of GUS expression for each
- 5 plasmid in transformed rice protoplasts, two wells from each thin layer culture were assayed 1, 3, 5, 10, 20 and 30 days after transformation. Cells were incubated with 5-bromo-4-chloro-3-indolyl glucuronide in a standard assay procedure for 48 hours before visualizing GUS activity.
- 10 For the quantitative analysis of GUS activity, cells were collected 20 days after transformation by low speed centrifugation in a bench top microcentrifuge. Total soluble protein was isolated in a GUS extraction buffer. Protein extracts
- 15 were incubated with 1 mM methylumbelliferyl- β -D-glucuronide (MUG) in a standard assay at 37°C for 3 to 6 hours. The liberation of 4-methylumbelliferone (MU) was followed by measuring fluorescence with excitation at 365 nm and emission at 455 nm in a spectrofluorometer. Protein concentrations of plant extracts were determined by the dye-binding method of Bradford.
- 20 Statistical analysis of the quantitative GUS activity results were done using a paired t-test.

To begin the analysis of the rice RAc1 regulatory regions, restriction mapping of a lambda-EMBL4 phage clone, lambda-RAc1, which had previously been shown to contain the rice RAc1 gene was carried out. The resulting restriction map of the 15.1 kb lambda-RAc1 insert is shown in Fig. 2A. This figure also indicates the position of the RAc1 coding and noncoding exons, as previously determined. To isolate and characterize the 5'-flanking sequence of the rice RAc1 gene, a 5.3 kb HindIII fragment from the lambda-RAc1 insert, spanning a region from 3.9 kb upstream of the RAc1 coding sequence to a point within its third coding exon, was isolated and cloned into the HindIII site of the plasmid pBluescript-KS to produce the plasmid pRAc15'.H3. A restriction map of the pRAc15'.H3 insert is shown in Fig. 2B. Restriction maps of the lambda-RAc1 (A) and pRAc15'.H3 (b) inserts were determined by single and double restriction enzyme digestion, as detailed in the methods. The enzyme sites are abbreviated as follows: BamHI, B; BglII, G; EcoRI, E; EcoRV, R; HincII, H2; HindIII, H3; SphI, Sp; SstI S; XhoI, Xh. The position of the 5.3 kb Hind III fragment within the lambda-RAc1 insert is indicated by dashed lines between (A) and (B). Noncoding and coding portions of RAc1 exons are depicted by open and filled boxes respectively. The strategy used to sequence the 2.1 kb EcoRV fragment within the pRAc15'.H3 insert is indicated by horizontal arrows. The length of the horizontal arrows are indicative of the size of the sequence obtained.

The restriction map of the pRAc15'.H3 insert shown in Fig. 2B also indicates the strategy used to sequence an EcoRV fragment which covers a region 2.1 kb upstream of the translation initiation codon of the RAc1 gene. This sequence is as follows:

					T
5	801	AGCTAGCATA	<u>CTCGAGGTCA</u>	TTCATATGCT	TGAGAAGAGA XhoI
		GTGCGGATAG	TCCAAAATAA	AACAAAGGTA	AGATTACCTG
	881	GTCAAAAGTG	AAAACATCAG	TTAAAGGTG	GTATAAGTAA
		AATATCGGTA	ATAAAAGGTG	GCCAAAGTG	AAATTACTCT
10	961	TTTTCTACTA	TTATAAAAA	TGAGGATGTT	TTTGCGGTAC
		TTTGATACGT	CATTTTGTGA	TGAATTGGTT	<u>TTTAAGTTTA</u>
	1041	TTGCGGATTT	TGGAAATGCA	TATCTGTATT	TGAGTCGGGT i
		<u>TTTAAGTTTG</u>	TTTGCTTTTG	TAAATACAGA	GGGATTGTGA i
	1121	TAAGAAATAT	CTTTAAAAA	AACCATATGC	TAATTGACA
15		TAATTTTGA	GAAAAATATA	TATTCAGGCG	<u>AATTCTCACA</u> Eco RI
	1201	ATGAACAATA	ATAAGATTAA	AATAGCTTGC	CCCCGTTGCA
		GCGATGGGTA	TTTTTTCTAG	TAAAATAAAA	GATAAACTTA
	1281	GACTCAAAAC	ATTTACAAAA	<u>ACAACCCCTA</u>	<u>AAGTCTTAAA</u> ii
		<u>GCCCAAAGTG</u>	<u>CTATGCACGA</u>	TCCATAGCAA	<u>GCCCAACCCA</u> iii
20	1361	<u>ACCCAACCGA</u>	<u>ACCCAACCGA</u>	<u>CCCCAGTGCA</u>	<u>GCCCACTGCG</u>
		AAATAGTCTC	CACACCCCGG	CACTATCACC	GTGAGTTGTG
	1441	CGCACCAACG	CAAGTCTGCG	<u>AGCCAAAAAA</u>	<u>AAAAAAGAA</u> iv
		<u>AGAAAAAAA</u>	<u>GAAAAAGAAA</u>	<u>AAACAGCAGG</u>	<u>TGGGTCCGGG</u>
	1521	TGCTGGGGGC	CGGAAAGCG	AGGAGGATCG	CGAGCAGCGA
25		CGAGCGCGGC	CCTCCCTCCG	CTTCCAAAGA	AAAGCCCCCC
	1601	ATCGGCACTA	<u>TATACATACC</u>	CCCCCTCTCT	CTCCCATCC
		<u>CCCAACCCCTA</u>	<u>CCACCAACAC</u>	<u>CACCAACACC</u>	<u>TCTCTCCCCC</u> v
	1681	<u>TGCTGCCCCG</u>	<u>ACGAAGAGCT</u>	<u>CCTCCCCCCCT</u>	<u>CCCCCTCCGC</u> vi
		<u>CGCCGCCGGT</u>	<u>AACCAACCCG</u>	<u>CCCTCTCTCT</u>	<u>CTTCTTTTCT</u> Sst I
30	1761	CGGTTTTTTT	TTTCGTCTCG	GTCTCGATCT	TTGGCCTTGG
		TAGTTTGGGT	GGGCGAGAGC	GGCTTCGTGG	CCAGATCCGG
	1841	TGGGCGGAG	GGGCGGGATC	TGGGCGCTGG	CGTCTCCGGG
		<u>CGTGAGTCGG</u>	<u>CCCGGATGCT</u>	<u>CGCGGGGAT</u>	<u>GGGCTCTCG</u> Bam HI

25

1921 GATGTAGATC TTCTTTCTTT CTTCITTTTG TGGGTAGAAT BglII
 TTGAATCCCT CAGCAITGTT CATGGTAGT TTTCTTTTC
 2001 ATGATTTGTG ACAAATGCAG CCTGGTGGG AGCTTTTTTG VII
 TAG GTAGAAAG ATG GCT GAC GCC GAG GAT ATC Eco RV
 5 Met Ala Asp Ala Glu Asp Ile

In this sequence, nucleotides are numbered with the A of the RAc1 transcription initiation site designated in bold. Restriction sites used in the subsequent construction of the various RAc1-GUS fusion plasmids and structural regions described in detail are underlined and named or designated by lower case Roman numerals and noted at the right margin of the sequence table. Upper case capital letters represent 5'-flanking sequences, upper case italic letters represent exon sequences and upper case capital letters between the two sets of exon sequences represent the intron sequence. The codons of the RAc1 first coding exon have their translation product indicated below them.

A number of potential regulatory sequences were identified in the 5'-flanking region of RAc1 gene. A 12 bp direct repeat, GGTTTTAAGTT (region i), is located between bases 1027 to 1038 and 1078 to 1088. A tandem (imperfect) direct repeat of 16 bp, AA(G/C)CCC(T)AAAGT(G/C)CTA (region ii), is located between bases 1301 and 1333. 20 bp downstream of this tandem direct repeat are eight tandem copies of an imperfectly repeating pentamer with the consensus sequence CCCAA (region iii). Finally, between bases 1465 and 1505, there is a purine rich sequence where 35 out of 40 bases are "A"s (region iv)

Downstream of these putative regulatory regions a number of sequences have been identified which have previously been implicated in the control of gene expression. Between bases 1609 and 1617 there is a putative TATA box (region v). The sequence and position of the putative RAc1 TATA box are in agreement with that previously determined for a number of other plant genes, and the sequence around the transcription initiation site; CCTACCA is similar to the consensus sequence for transcription initiation previously determined for a number of animal YAYY and plant YYYAYA genes (Y= pyrimidine). The noncoding exon located 3' of the TATA box is GC rich (77.5%) and consists of a number of tandemly repeated (A/T)CC triplets (region vi).

The 5'-intron of the rice RAc1 gene is 313 bp long. Its 5'-acceptor splice site [(G/G)TA], and 3'-donor splice site [TTTTTTGTA(G/G)], follow the consensus sequence previously determined for a number of plant genes. A putative branch point site for mRNA splicing between bases 2008 and 2012 (region vii) was identified whose sequence, GTGAC, and distance from the 3'-donor splice site bears similarity to the location and consensus sequence for animal mRNA branch point splice sites, YTRAC.

Four RAc1-GUS fusion plasmids were constructed to determine the minimum amount of RAc1 5'-flanking sequence required for maximal β -glucuronidase (GUS) gene expression; these are shown in Fig. 3. The plasmid pRAc15'.21VG has the 2.1 kb EcoRV restriction fragment from pRAc15'.H3 fused to the GUS coding region and 3'-noncoding transcript terminator region of the

nopaline synthase (NOS) gene. This plasmid encodes a transcript containing two in frame translation start codons, one each from the RAc1 and GUS genes, adding 15 amino acids to the N-terminal end of the wild type GUS protein. The plasmids pRAc15'.13XG and
5 pRAc15'.09RG were created by the deletion of 0.8 and 1.2 kb of sequence, respectively, from the 5' end of pRAc15'.21VG. The plasmid pRAc15'.21VR'G has the 5' 1.2 kb EcoRI fragment cloned in the opposite orientation to that in pRAc15'.21VG.

A number of plasmids were also constructed to investigate
10 the importance of the RAc1 5'-intron on RAc1-promoter-mediated gene expression. The plasmid pRAc15'.09RGIΔ8 contains a deletion of 9 bp around the BglII site of the RAc1 5'-intron. The plasmid pRAc15'.09RGIΔ1 contains a deletion of 133 bp between bases 1868 and 2003 of the sequence depicted above, reducing the
15 distance between the 5' acceptor and mRNA branch point from its 5'-intron. The plasmid pRAc15'.09RGIΔ12 has a deletion of 157 bp between bases 1868 and 2027, removing the putative mRNA branch point from its 5'-intron. The plasmid pRAc15'.09RGIDS- was created by excision of a 170 bp BamHI fragment from
20 pRAc15'.09RG, removing the putative mRNA branch point and 3'-donor splicing sites from the RAc1 5'-intron. The plasmid pRAc15'.09RGI- is identical to pRAc15'.09RG except that it lacks the entire RAc1 5'-intron.

The plasmid pRAc15'.21V'G contains the 2.1 kb EcoRV
25 fragment from pRAc15'.H3 cloned in the opposite orientation to that in pRAc15'.21VG. It was postulated that the RAc1 5'-region

would not show bidirectional promoter activity because the 5'-most 1.2 kb EcoRV-EcoRI restriction fragment in pRac15.21VG did not show any binding when used as a probe in Northern hybridization against total RNA from 7 day old rice shoots. As a positive control, the plasmid pAl₁GusN was used. This plasmid contains the promoter, first exon and first intron of the maize Adh1 gene fused to a GUS coding sequence with the 3' noncoding region of the NOS gene.

More specifically, the individual maps of constructs containing various portions of the 5'-flanking and 5'-transcribed sequence of the rice Rac1 gene fused, in frame, to a sequence containing the GUS coding region and 3' NOS transcription terminator are shown in Figure 3. Open boxes represent noncoding exons of the Rac1 gene, filled boxed represent the GUS coding region (not to scale) and striped boxes represent the NOS terminator sequence. The stippled box in the construct pAl₁GusN represents the first exon of the maize Adh1 gene. The 'Δ' symbol indicates the deletion point in the various Rac1 first intron deletion constructs. The indicated restriction enzyme sites are abbreviated as follows: BamHI, B1; BglII, B2; EcoRI, R1; EcoRV, RV; HincII, H2; HindIII, H3; SstI, S1; XhoI, Xh1; XbaI, Xb1.

To assay GUS expression from the various Rac1-GUS fusion plasmids, rice suspension culture cells were transformed by the biolistic method [see Wang, Y.C. et al., Transient expression of foreign genes in rice, wheat, and soybean cells following particle bombardment. Plant Mol. Biol. 11:433 (1987); and Cao, J. et al.,

Transformation of rice and maize using the biolistic process. UCLA Symposia on Plant Gene Transfer (1989)], and assayed for high level GUS activity by visual inspection 10 days after transformation according to Jefferson et al. [GUS fusions: β -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO J. 6:3901 (1987)].

The results of these various assays are tabulated in the following table:

TABLE I

	Protoplasts:		Cell Suspension	
	Names	mean specific Activity nmol/hour/ μ g	mean GUS Expression %	Visible GUS Activity
10				
15	pRAc15'.21VG	211.4 ¹	8.4	+
	pRAc15'.13XG	195.0 ¹	7.9	+
	pRAc15'.09RG	94.2 ²	5.7	+
	pRAc15'.21VR'G	92.5 ²	6.1	+
20	pRAc15'.09RGI Δ 8	95.4 ²	2.8	N.A.
	pRAc15'.09RGI Δ 1	53.4 ³	1.4	N.A.
	pRAc15'.09RGI Δ 12	17.0 ⁴	0.2	N.A.
	pRAc15'.09RGIDS-	8.8 ⁵	0.2	N.A.
	pRAc15'.09RGI-	4.7 ⁵	0.2	-
25	pRAc15'.21V'G	0.7 ⁵	0.0	-
	Untransformed	0.0 ⁵	0.0	-
	pAl ₁ GusN	31.9 ⁶	4.2	+

N.A. = not assayed

The results of the rapid assay are shown in the third column of Table 1. Suspension cultures transformed with the plasmids pRac15'.21VG, pRac15'.13XG, pRac15'.09RG, pRac15'.21VR'G and pAl₁GusN were all positive for GUS activity while untransformed cultures or those transformed with pRac15'.09RGI⁻ or pRac15'.21V'G displayed no visible GUS expression.

Table I also shows the results of a determination of GUS expression efficiencies, defined as the percent of intact cells displaying visible GUS activity 10 days after PEG-mediated transformation of rice protoplasts with the Rac1-GUS fusion constructs. The plasmids pRac15'.21VG and pRac15'.13XG displayed the highest GUS expression efficiencies at 8.4 and 7.9% respectively. The plasmids pRac15'.09RG and pRac15'.21VR'G showed GUS expression efficiencies of 5.7 and 6.1%, respectively.

All four of the aforementioned plasmids displayed significantly higher GUS expression efficiencies than was found for pAl₁GusN. Removal of Rac1 5'-intron sequences resulted in a significant reduction in the number of cells displaying visible GUS activity, relative to pRac15'.09RG, which contains the intact Rac1 5'-intron. For plasmids pRac15'.09RGIΔ8 (2.8%) and pRac15'.09RGIΔ1 (1.4%), the longer the sequence deleted between the 5'-acceptor and the putative mRNA branch point splice sites, the lower the GUS expression efficiency. Removal of the branch point for mRNA splicing in the plasmid pRac15'.09RGIΔ12 and the additional removal of the 3'-donor splice site in the plasmid pRac15'.09RGIDS⁻ led to GUS expression efficiencies which, at

0.2%, represents less than 4% of that determined for the plasmid pRac15'.09RG. Untransformed protoplasts or protoplasts transformed with either pRac15'.21V'G or the intronless plasmid pRac15'.09RG1- gave no blue cells.

- 5 To determine the quantitative differences in GUS expression from the various Rac1-GUS fusion plasmids, the constructs were introduced into rice protoplasts by PEG-mediated transformation, and GUS specific activity was assayed 20 days after the transformation procedure. The results of this quantitative
10 analysis of GUS expression are presented in the first column of Table I.

- The highest GUS specific activities were recorded for the plasmids pRac15'.21VG and pRac15'.13XG, which have the longest Rac1 upstream regions. The plasmid pAl₁GusN, which has the
15 GUS gene expresses from the maize Adh1 promoter, displayed less than 17% of the GU specific activity shown by pRac15'.21VG and pRac15'.13XG. The plasmids pRac15'.09RG, with a truncated Rac1 5'-region, and pRac15'.21VR'G, with an inverted Rac1 5'-region, were found to have GUS specific activities around 48% of that for
20 pRac15'.21VG. There was no statistically significant difference (mean specific activity with identical superscripted numbers are not statistically different from each other), in specific activity between these two constructs, suggesting that the inversion of the Rac1 5'-end did not influence GUS expression from the
25 truncated Rac15'.09RG construct.

The effects of the various RAc1 intron deletions could be seen when their GUS specific activities were compared to that of their progenitor plasmid, pRAc15'.09RG. The small 9 bp deletion in the RAc1 5'-intron of plasmid pRAc15'.09RGIΔ8 had no effect on the quantitative level of GUS activity. The plasmid pRAc15'.09RGIΔ1, containing a deletion of 133 bp between the 5'-acceptor splice site and the putative mRNA branch point splice site, displayed a GUS specific activity that was less than 57% of that observed for pRAc15'.09RG. The plasmid pRAc15'.09RGIΔ12, which had the putative mRNA branch point site removed, showed a further decrease in GUS specific activity of 17% of that observed for pRAc15'.09RG. The plasmid pRAc15'.09RGIDS⁻, which lacks the putative mRNA splicing branch site and 3'-splicing donor site of the RAc1 5'-intron, and pRAc15'.09RGI⁻, which lacks the entire RAc1 5'-intron, displayed no significant GUS specific activities over that observed for untransformed samples. However, it was noted in Table I that while blue cells, indicative of GUS expression, were never observed in untransformed rice protoplasts or protoplasts transformed with the intronless plasmid pRAc15'.09RGI⁻, GUS expression was able to be visualized in those rice protoplast transformed with the plasmid pRAc15'.09RGIDS⁻.

As a first step towards an understanding of the regulation of rice actin gene expression, the present invention has described the isolation and characterization of the 5'-regions of the rice RAc1 gene. Within the 5'-flanking sequence, the occurrence of a

long poly(dA) stretch located between bases 1465 and 1505 which appears to play a role in the constitutive activation of RAC1 gene expression was found. Based upon chemical analysis, it has been suggested that a minimal affinity for histone cores and
5 nucleosome formation is provided by homogeneous tracts of purines located on one strand of the double helix. Poly(dA) regions within recombinant DNA molecules can prevent nucleosome formation *in vitro*, and it has also been found that naturally occurring poly(dA) tracts act as 5'-promoter elements
10 for the constitutive expression of different yeast genes. It is believed that poly(dA) stretches may change the chromatin structure, enabling general transcription factors to access the DNA template and activate constitutive transcription in the absence of more specific transcription factors.

15 The sequence of RAC1 untranslated mRNA 5'-region is unusually GC rich, with an AT content of only 26.5%. In a survey of 79 plant genes it was found that their mRNA 5'-sequences were extremely AT rich, with 71 of the genes surveyed having an AT content greater than 51% and only one of the genes surveyed
20 having an AT content of less than 44%. In the ribosomal-scanning model of translation initiation it is proposed that the AT richness of the mRNA 5'-sequence leads to the formation of relatively labile secondary structures which facilitate the movement of the ribosomal subunits towards the translation initiation codon. The
25 significance of the GC rich RAC1 mRNA 5'-sequence, with respect

to transcript stability and translation efficiency, remains to be investigated.

The construction of a number of RAc1-GUS fusion constructs has provided the determination that the plasmid pRAc15'.13XG, containing 0.83 kb of 5'-flanking sequence, the noncoding exon and 5'-intron of the RAc1 gene, has the minimal amount of RAc1 sequence necessary for maximal GUS expression in transient assays of transformed rice protoplasts; the additional 0.8 kb of RAc1 5'-sequence presents in the pRAc15'.13VG did not significantly increase GUS expression above that of the plasmid pRAc15'.13XG. It was also found in the present invention that the RAc1 5'-flanking sequence was more active than the maize Adh1 5'-flanking region in stimulating GUS expression in transformed rice protoplasts. Deletion of a 0.4 kb region from the 5'-end of the pRAc15'.13XG insert resulted in a 52% reduction in GUS activity. However, no statistically significant difference in GUS specific activities was detected between the plasmids pRAc15'.09RG, with the 0.4 kb region deleted, and pRAc15'.13.09RG, with the 0.4 kb deleted, and pRAc15'.21VR'G, with the 0.4 kb region present but inverted and displaced 0.8 kb further upstream. This suggests that the sequence element(s) in the 5'-end of the pRAc15'.13XG insert that are responsible for high level GUS expression do not display any position/orientation-independent, enhancer-like activity. In most cases the differences in GUS specific activities detected between the RAc1-GUS fusion constructs could be directly correlated with

their differential GUS expression efficiencies. This suggests that there is a threshold to the visualization of GUS expression. Protoplasts transformed with the various RAc1-GUS fusion constructs are presumed to display a position-effect-dependent distribution of GUS specific activities, the mean of which is a function of their particular RAc1 promoter. Therefore those protoplast populations transformed with the most active RAc1 promoter-GUS fusion constructs will display the greatest number of visually detectable blue cells and the highest GUS expression efficiencies.

By constructing RAc1-intron-deletion-GUS plasmids, it was able to be shown that GUS expression in transformed rice protoplasts was dependent on the presence of an intact RAc1 5'-intron. Deletion of the RAc1 intron reduced GUS specific activity to levels that were not significantly greater than that of untransformed protoplasts. In transient assays of transformed maize protoplasts a maize Adh1-S gene lacking the nine Adh1-S introns was expressed at levels 50- to 100-times lower than that of the intact gene. The reintroduction of the 5'-intron of the Adh1-S gene was enough to restore the level of expression to that observed for the intact gene, although this effect was only observed if the first intron was reinserted in a 5'-position; it is believed that this was a result of increases in the amount of mature, cytoplasmic mRNA and not a result of increased transcript stability in the nucleus or increased translation efficiency in the cytoplasm.

The increase in expression resulting from the presence of RAc1 5'-intron is not believed to be caused by the presence of some enhancer-like cis acting sequence within the intron. No sequences within the RAc1 intron were found which bore any homology to either an enhancer-like sequence from the first intron of the human β -cytoplasmic actin gene, or to any sequences within the maize Adh1 5'-intron. Nor is it believed that the RAc1 first intron codes for any functionally active transcript or protein product. Previous Northern blot hybridization with a double stranded probe that spans the RAc1 first intron failed to show any binding to total RNA from seven day old rice shoots, other than to the 1.7 kb RAc1 transcript itself. Finally, the RAc1 5'-intron contains no open reading frames of any significant length. One conclusion of the present invention suggests that the primary effect of RAc1 first intron is associated with an *in vivo* requirement for efficient splicing.

The small deletion in the RAc1 intron in the plasmid pRAc15'.09RGI Δ 8 caused no significant reduction in GUS specific activity relative to that of pRAc15'.09RG. However, a significant difference was observed between the GUS specific activities of the plasmids pRAc15'.09RG and pRAc15'.09RGI Δ 1. The 133 bp deletion in the plasmid pRAc15'.09RGI Δ 1, while not removing any of the sequences previously implicated in intron splicing, did cause a reduction in the distance between the 5'-acceptor and putative mRNA branch point splice sites with an associated 44% reduction in GUS specific activity. Large reductions in splicing

efficiency occur *in vivo* and *in vitro* when the distance between the 5'-acceptor and branch point splice sites is reduced; this requirement for a minimal distance between the 5'-acceptor and branch point splice sites probably reflects a requirement for multiple splicing factors to interact with specific intron regions, these regions being presumably removed in the construction of pRAC15'.09RGIΔ1. The *in vivo* requirement for efficient intron splicing was further supported by the observation that a deletion which removed the putative 5'-intron mRNA branch point splice site in the construction of pRAC15'.09RGIΔ12, or removed both the branch point and 3'-donor splice sites in the construction of pRAC15'.09RIDS⁻ led to GUS specific activities which were 18% and 9% of that observed for pRAC15'.09RG. Deletion of the normal branch point and 3'-donor splice sites of animal genes does not abolish splicing but rather leads to the use of cryptic branch point and 3'-donor splice sites, with associated reductions in splicing efficiency and gene expression; this may be occurring in rice protoplasts transformed with the plasmids pRAC15'.09RGIΔ12 and pRAC15'.09RGIDS⁻, leading to reduced but still detectable GUS expression

The results of the present invention clearly demonstrate that the RAc1 5'-flanking sequence between 0 and 2070 nucleotides and more particularly between 800 and 2070 nucleotides in the genomic sequence contains an efficient promoter for rice transformation. Furthermore, the expression of a foreign gene in transformed plant cells can be dependent upon,

but not necessarily an absolute function of, the presence of an intact 5'-intron sequence. It appears that a functional requirement for the presence of the 5'-intron may be correlated with the conservation of the 5' noncoding exon during this rice actin gene's structural evolution. In addition the results of the present invention indicate that the maize 5'-intron is located between the first and second coding exons of the Adh1 gene, while the rice 5'-intron is located between a 5' noncoding exon and the first coding exon of the RAc1 gene, suggesting that there may be a common positional component to the intron mediated stimulation of gene expression observed for the maize and rice 5'-intron.

In summary, the preceding description of the present invention clearly demonstrates that a 2.1 kbp 5' of the Act1 gene's translation initiation codon, containing 1.3 kb of 5' untranslated sequence, the 5' transcribed but untranslated exon, 5'-intron and part of the first coding exon of the rice Act1 gene, is capable of conferring high level expression of foreign gene in transformed rice material. Thus this region can be used to activate the constitutive expression of foreign genes in transgenic plants of rice and other agronomically important plants; the 5'-intron of the rice Act1 gene can stimulate the expression of a foreign gene in transformed rice material [thus this (and the other introns of the rice Act1 gene) will be able to increase the expression of foreign genes in transformed plants of rice (and other agronomically important plants) when ligated in front of such a gene in between a promoter and the gene]; and

although no specific function for the 3'-end of the rice Act1 gene in the regulation of foreign gene expression in rice has been demonstrated. It appears from the present invention that the 3'-end of the rice Act1 gene should also stimulate the expression of such foreign genes in transformed plants of rice and other agronomically important crops.

Thus, while we have illustrated and described the preferred embodiment of our invention, it is to be understood that this invention is capable of variation and modification and we therefore do not wish or intend ourselves to be limited to the precise terms set forth, but desire to avail ourselves of such changes and alterations which may be made for adapting the invention to various usages and conditions. Accordingly, such changes and alterations are properly intended to be within the full range of equivalents, and therefore within the purview of the following claims.


Having thus described our invention and the manner and a process of making and using it in such clear, full, concise and exact terms so as to enable any person skilled in the art to which it pertains, or with which it is mostly nearly connected, to make and use the same;

WE CLAIM:



1. A promoter for monocotyledonous plants including a DNA sequence comprising a 5'-region of a rice RAc1 gene that directs the expression of a linked structural gene in the tissues of a recipient plant.
2. A promoter for monocotyledonous plants according to Claim 1 comprising the region extending from approximately position 1 to approximately position 2070 of the RAc1 gene.
3. A promoter for monocotyledonous plants according to Claim 2 containing the region extending from approximately position 800 to approximately position 2070 of the RAc1 gene.
4. A recombinant monocotyledonous plant comprising a structural gene for the production of a plant produced product linked with a promoter comprising a 5'-region of a rice RAc1 gene that directs the expression of the linked structural gene.
5. A genetic construct selected from the group consisting of pRAc15'.21VG, pRAc15'.13XG, pRAc15'.09RG, pRAc15'.21VR'G, pRAc15'.09RGIΔ8, and pRAc15'.09RGIΔ1.
6. A regulatory element for RAc1 gene promoter activity comprising approximately a 1.3 kb genetic sequence located in the 5'-region upstream portion of the RAc1 translation initiation codon.
7. A genetic construct comprising the RAc15' intron ligated in between a suitable promoter and a structural gene to enhance the expression of the latter.

8. A genomic sequence comprising the RAc1 sequence including a 5' non-coding exon separated by a 5'-intron from the first coding exon in the sequence.

9. A genomic sequence comprising a fragment having a restriction map selected from:




1/2

FIG. 1**FIG. 2**

SUBSTITUTE SHEET

2/2

FIG. 3**SUBSTITUTE SHEET**

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/00073

I. CLASSIFICATION * SUBJECT MATTER (in several classification symbols, none)

According to International Patent Classification (IPC) or to both National Classification and IPC
 ICP (5): C12N 15/29, 15/63, 15/67, 15/82; A01H 4/00, 5/00
 U.S. Class: 435/172.3; 536/27; 800/205

II. FIELDS SEARCHED

Minimum Documentation Searched *

Classification System	Classification Symbols
U.S.	435/172.3, 317.1; 536/27; 935/30, 35; 800/205, Dig. 52, Dig. 57

Documentation Searched other than Minimum Documentation
 to the extent that such Documents are included in the Fields Searched *

Chemical Abstracts on line (File Biosis, 1969-1991); USPTO automated patent system
 (file USPAT, 1971-1991); sequence search (Gene bank and URMEL). See attachment for search
 terms.

III. DOCUMENTS CONSIDERED TO BE RELEVANT *

Category *	Citation of Document, II with indication, where appropriate, of the relevant passages in	Relevant to Claim No. 1
X.P.	Plant Molecular Biology, volume 14.	1-3, 6-9
Y.P.	issued 1990. McElroy et al. "Structural Characterization of a rice actin gene". pages 163-171. see the entire document..	4-5
X.P.	The Plant Cell, volume 2, issued February	1-3, 6-9
Y.P.	1990. McElroy et al. "Isolation of an efficient actin promoter for use in rice transformation." pages 163-171. see the entire document.	4-5
Y	Genes and Development, volume 1, issued 1987. Callis et al. "Introns increase gene expression in cultured maize cells," pages 1183-1200. see the entire document.	1-7

* Special Categories of cited documents: (1)

"A" document claiming to be a material state of the art which is not
considered by the International Searching Authority

"E" earlier document but published on or after the international
filing date

"L" document which is a prior art document, published on or
after the international filing date, which is not a material state of the art
which is not considered by the International Searching Authority

"O" document relating to the same subject matter as the international
application, but not published on or after the international filing date

"P" document which is a prior art document, published on or
after the international filing date, which is not a material state of the art
which is not considered by the International Searching Authority

IV. CERTIFICATION

Date of the International Search Report: 28 March 1991

28 March 1991

International Searcher's Name: ISA/US

ISA/US

09 MAY 1991

L. McElroy
 Che S. Chereskin

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	Plant Cell Reports, volume 8, issued 1989. Oard et al. "Chimeric gene expression using maize intron in cultured cells of breadwheat" pages 156-160. see the entire document.	1-7,
Y	Gene, volume 65, issued 1988. Nairn et al. "Nucleotide sequence of an actin gene from <u>Arabidopsis Thaliana</u> ." pages 247-257, see the entire document, but especially Figure 4 and page 248. column 1.	1-3,5-9
Y	Nucleic Acids Research, volume 17, number 1, issued 1989. Frederickson, et al. "5' Flanking and first intron sequences of the human β -actin gene required for efficient promoter activity." pages 253-270. see the entire document, but especially pages 254 and 261-263.	1-3,5-9
Y	DNA, volume 2, number 1, issued 1981. Nagao et al. "Multigene family of actin-related sequences isolated from a soybean genomic library". pages 1-9 see the entire document.	1-9
Y	Nuclear Acids Research, Volume 17, number 2, issued 1989. Ng et al. "Regulation of the human β -actin promoter by upstream and intron domains". pages 601-615. see the entire document.	1-7
Y	Journal of Molecular and Applied Genetics, volume 2, No. 1 issued 1983, Shah et al. "Genes encoding actin a higher plants: intron positions are highly conserved but the coding sequences are not." pages 111-126. see the entire document.	1-9
A	Protoplasma, volume 141, issued 1987. Staiger et al: "Actin localization and function in higher plants." pages 1-12. see the entire document.	1-9

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☐ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out², specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING³

This International Searching Authority found multiple inventions in this international application as follows:

- I. Claims 1-3, 5-7 drawn to promoter and regulatory elements.
- II. Claim 4 drawn to recombinant plants.
- III. Claims 8-9 drawn to genomic sequences.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report covers only the invention first mentioned in the claims; if it is covered by claim number(s):

4. ☐ As all search fees have been timely paid by the applicant, this international search report covers all searchable claims of the international application.

Remarks on Prior Art:

- ☐ The claims of the international application are not drawn to a single general inventive concept.
- ☐ No request for amendment of the international application was filed.